arXiv:1207.0779v1 [hep-ph] 3 Jul 2012

Flavor-Tuned SUSY Higgs Bosons at the LHC: Enhanced Di-photon and Suppressed Di-tau Signals of the 125 GeV Higgs

Vernon Barger*

Department of Physics, University of Wisconsin, Madison, WI 53706, USA

Muneyuki Ishida†

Department of Physics, Meisei University, Hino, Tokyo 191-8506, Japan

Wai-Yee Keung‡

Department of Physics, University of Illinois at Chicago, IL 60607, USA

(Dated: July 4, 2012)

Abstract

We explain the enhanced di-photon signal of the 125 GeV Higgs boson, h, observed by the ATLAS and CMS collaborations by identifying it with the neutral H_u^0 of the two Higgs doublets of minimal Supersymmetry. The off-diagonal element of the (H_u^0,H_d^0) mass matrix in the flavor basis must be suppressed so that there is nearly no mixing. A direct corollary is the suppression of h to di-tau and b-quark pair decays. The constraints on the other SUSY Higgs states (H, A, H^+) are severe; all must have masses comparable to m_h and their LHC signals are unambiguously predicted.

A $\gamma\gamma$ enhancement of the 125 GeV Higgs boson signal relative to the Standard Model (SM) Higgs expectation has been reported by the ATLAS[1] and CMS[2] experiments at the LHC. We explain this result in minimal Supersymmetry by a nearly zero mixing between the neutral CP-even Higgs flavor states H_u^0,H_d^0, with the 125 GeV Higgs signal identified with H_u^0 . The 125 GeV Higgs to $\tau\tau$ decay is correspondingly predicted to be almost absent. The required suppression of the off-diagonal element of the (H_u^0,H_d^0) mass matrix in the flavor basis implies a cancellation of tree-level and loop-contributions and implies a light mass spectrum of the the other Higgs bosons (H^0, A^0, H^+). The other CP-even Higgs, H^0, is identified with the flavor basis state H_d^0. The production and decays at the LHC of these yet-to-be observed Higgs SUSY Higgs bosons are predicted.

Ratios of the SUSY Higgs couplings to those of the SM Higgs The SUSY Higgs mechanism is based on the two Higgs doublet model of type II[3–5] with one Higgs doublet H_u coupled to up-type quarks and one Higgs doublet coupled to down-type quarks. After spontaneous symmetry breaking the physical Higgs states are two CP-even neutral Higgs h, H, one CP-odd neutral pseudo-scalar A and a charged Higgs $H^{\pm}$. The neutral Higgs bosons are related to the flavor eigenstates H_u^0 and H_d^0 by

$$\begin{pmatrix} H_u^0 \\ H_d^0 \end{pmatrix} = \begin{pmatrix} v_u \\ v_d \end{pmatrix} + \frac{1}{\sqrt{2}} \begin{pmatrix} c_\alpha & s_\alpha \\ -s_\alpha & c_\alpha \end{pmatrix} \begin{pmatrix} h \\ H \end{pmatrix} + \frac{i}{\sqrt{2}} \begin{pmatrix} s_{\beta^0} & c_{\beta^0} \\ -c_{\beta_0} & s_{\beta_0} \end{pmatrix} \begin{pmatrix} G \\ A \end{pmatrix} \tag{1}$$

where α is the CP-even Higgs-mixing angle, and β_0 is the CP-odd mixing angle which coincides with β of tan$\beta \equiv v_u/v_d$. G is Nambu-Goldstone boson that becomes the longitudinal degree of the Z boson. We use the notation $s_\alpha = \sin\alpha$, $c_\alpha = \cos\alpha$, and $t_\alpha = \tan\alpha$.

We focus first on the lightest Higgs boson h, which has the composition

$$\frac{h}{\sqrt{2}} = c_\alpha H_u^0 - s_\alpha H_d^0 \ . \tag{2}$$

The ratios of the h couplings to those of the SM Higgs h_{SM}, which are denoted as $r_{CC}^h (\equiv g_{hCC}/g_{h_{SM}CC})$, are given by

$$r_{VV}^h = \sin(\beta - \alpha), \qquad r_{tt}^h = r_{cc}^t = \frac{c_\alpha}{s_\beta}, \qquad r_{bb}^h = r_{\tau\tau}^h = r_{\mu\mu}^h = \frac{-s_\alpha}{c_\beta} \ . \tag{3}$$

The corresponding partial decay widths relative to those of h_{SM} are given by the absolute square of these r_{CC}^h.

By using these coupling ratios, the $gg, \gamma\gamma$ coupling ratios $r^h_{gg,\gamma\gamma}$ to those of h_{SM} are obtained[6] by

$$r^h_{gg} = \frac{1.03\frac{c_\alpha}{s_\beta} + (-0.059 + i0.081)\frac{-s_\alpha}{c_\beta}}{1.03 + (-0.059 + i0.081)} \ , \tag{4}$$

$$r^h_{\gamma\gamma} = \frac{\frac{7}{4}1.19 \ \sin(\beta - \alpha) - \frac{4}{9}1.03\frac{c_\alpha}{s_\beta} - \frac{1}{9}(-0.059 + i0.081)\frac{-s_\alpha}{c_\beta}}{\frac{7}{4}1.19 - \frac{4}{9}1.03 - \frac{1}{9}(-0.059 + i0.081)} \ , \tag{5}$$

where the first(second) terms in the numerator and denominator in Eq. (6) represent the top(bottom)-quark triangle-loop contributions, and the first terms in the numerator and denominator in Eq. (7) represent the W-boson triangle-loop contribution. The bottom quark contribution in r^h_{gg} becomes significant compared with the top contribution only when tan$\beta \sim 10$.

The cross section ratios[6] of the process, $XX \to h \to CC$ relative to h_{SM} are obtained as follows

$$XC \equiv \frac{\sigma_{XX\to CC}}{\sigma_{XX\to h_{SM} \to CC}} = \frac{|r^h_{XX} r^h_{CC}|^2}{R^h} \ , \tag{6}$$

$$R^h = 0.58|r^h_{bb}|^2 + 0.06|r^h_{\tau\tau}|^2 + 0.24|r^h_{VV}|^2 + 0.09|r^h_{gg}|^2 + 0.03|r^h_{cc}|^2 \ , \tag{7}$$

where R^h is the ratio of the h total width to that of h_{SM} and we have used the branching fractions of the 125 GeV h_{SM} decays to $b\bar{b}, \tau\tau, VV(= WW^* + ZZ^*), gg, c\bar{c}$, which are predicted[7], respectively, as 0.58, 0.06, 0.24, 0.09, 0.03. XC for the various processes are determined only by two parameters, α and tanβ. They are plotted as a function of α in Fig. 1 for tan$\beta = 10$.

It can be seen in Fig. 1 that the $\gamma\gamma$ cross sections relative to h_{SM} are sharply enhanced and reach a maximum of about 3[8] in the vicinity of $\alpha = 0$, where $h = H^0_u$. There, the $\tau\tau$ and b-pair decays are highly suppressed. The $\tau\tau$ suppression factor is

$$|r^h_{\tau\tau}| = \left|\frac{-s_\alpha}{c_\beta}\right| \ll 1 \to |\alpha| \ll \frac{1}{\tan\beta} \tag{8}$$

for moderate values of tanβ, tan$\beta > 5$. Thus, α must be flavor-tuned to be close to zero in order to reproduce the $\gamma\gamma$ enhancement. This flavor-tuned Higgs boson is different from the decoupling limit where $\alpha = \beta - \frac{\pi}{2}$ and $r^h_{\tau\tau} = 1$.

The predicted values of XC for $\alpha = 0$ are compared with the experimental data in Table I. The maximum value of $g\gamma = 3.1$, but a small non-zero α value can lead to $g\gamma \sim 2$. Correspondingly, $gb \sim 0.6$ The XC are almost independent of the value of tanβ. The



FIG. 1. α dependence of the cross-sections XC of various processes relative to the SM Higgs. We consider five quantities $XC = g\gamma, V\gamma, gV,\ Vb$ and $g\tau$, corresponding to $gg \to \gamma\gamma$(dotted black), $VV \to \gamma\gamma$(dashed green), $gg \to VV(WW^*$ or $ZZ^*)$(long-dashed blue) which almost overlaps with $g\gamma$, $gg \to b\bar{b}, \tau\bar{\tau}, \mu\bar{\mu}$(solid orange), and $VV \to b\bar{b}, \tau\bar{\tau}$(dot-dashed red). $h = H_u^0(H_d^0)$ and $H = H_d^0(H_u^0)$ when $\alpha = 0(\pi/2)$. Here the $\alpha = \pi/2$ corresponds to $H = H_d^0$ with the same mass.

predicted suppression of the b-pair signal is at variance with the reported Tevatron[11] signal which has about a 3 sigma significance. We predict $g\gamma = gZ$. Thus, the actual ratio of the event numbers is calculated as $\sigma_{gg\to ZZ^*\to 4l}/\sigma_{gg\to\gamma\gamma} = BF(Z \to l^+l^-)^2 BF(h_{SM} \to ZZ^*)/BF(h_{SM} \to \gamma\gamma) = (0.0337)^2 \times 0.0264/0.00228 = 0.0131$.

By using the coupling ratios r_{CC}^h with $\alpha = 0$, the h total width[6] is obtained as $\Gamma_{\rm tot}^h = 1.5$ MeV which is compared with $\Gamma_{\rm tot}^{h_{SM}} = 4.07$ MeV[7] at $m_{h_{SM}} = 125$ GeV.

Flavor-Tuning of the mixing angle α The squared-mass matrix of the neutral Higgs in the u, d basis is given in the 1-loop level[9, 10] approximation as

$$\mathcal{M}_H^2 = \begin{pmatrix} m_A^2 c_\beta^2 + M_Z^2 s_\beta^2 + {\rm Loop}_{11} & -(m_A^2 + M_Z^2) s_\beta c_\beta + {\rm Loop}_{12} \\ -(m_A^2 + M_Z^2) s_\beta c_\beta + {\rm Loop}_{12} & m_A^2 s_\beta^2 + M_Z^2 c_\beta^2 \end{pmatrix} ,$$

$$\mathrm{Loop}_{12} = \frac{m_t^4}{16\pi^2 v^2 s_\beta^2} \frac{\mu X_t}{M_{\rm SUSY}^2} \left[\frac{A_t X_t}{M_{\rm SUSY}^2} - 6 \right] + \frac{h_b^4 v^2}{16\pi^2} s_\beta^2 \frac{\mu^3 A_b}{M_{\rm SUSY}^4}$$

$$\mathcal{M}_H^2 \to \begin{pmatrix} M_Z^2 + {\rm Loop}_{11} & -\epsilon + {\rm Loop}_{12} \\ -\epsilon + {\rm Loop}_{12} & m_A^2 \end{pmatrix} ,$$

$$\mathrm{Loop}_{12} \to \frac{m_t^4}{16\pi^2 v^2} \frac{\mu X_t}{M_{\rm SUSY}^2} \left[\frac{A_t X_t}{M_{\rm SUSY}^2} - 6 \right] + \frac{h_b^4 v^2}{16\pi^2} \frac{\mu^3 A_b}{M_{\rm SUSY}^4}, \tag{9}$$

	$\sigma_h/\sigma_{h_{\rm SM}}$	FT($\alpha = 0$)	($\alpha = 0.06$)	CMS[2]	ATLAS[1]	Tevatron[11]
$VV \to \tau\tau$	$V\tau$	0	0.60			
$q\bar{q} \to Vb\bar{b}$	qb	0	0.60	$1.2^{+2.1}_{-1.9}$	$-0.8^{+1.8}_{-1.7}$	2.0±0.7
$gg \to \tau^-\tau^+$	$g\tau$	0	0.74	$0.63^{+1.00}_{-1.28}$	0.0±1.7	
$gg \to \gamma\gamma$	$g\gamma$	3.1	1.9	1.62±0.68	$1.6^{+0.8}_{-0.7}$	$3.4^{+3.1}_{-2.4}$
$gg \to WW^*$	gW	3.1	2.0	0.40±0.55	0.20±0.62	$0.0^{+1.0}_{-0.0}$
$gg \to ZZ^*$	gZ	3.1	2.0	$0.58^{+0.94}_{-0.58}$	$1.4^{+1.3}_{-0.8}$	
$VV \to \gamma\gamma$	$V\gamma$	2.7	1.6	$3.8^{+2.4}_{-1.8}$[2]		

TABLE I. $XC \equiv \sigma(X\bar{X} \to C\bar{C})/\sigma(X\bar{X} \to h^0_{\rm SM} \to C\bar{C})$: The Higgs-signal cross sections at the LHC relative to the SM Higgs cross sections in the flavor-tuned (FT) model with $\alpha = 0$ and $\alpha = 0.06$ in the case tan$\beta = 10$. These values are compared with the experimental values by CMS[2] and by ATLAS[1]. $VV \to \tau\tau, \gamma\gamma$ are determined by CMS from di-jet diphoton events. The $b\bar{b}$ signal of the second row is inferred from the Tevatron data[11]. The cross sections are obtained from $XC \times \sigma_{XX \to h_{SM}} BF(h_{SM \to CC})$ where the h_{SM} production cross sections $\sigma_{XX \to h_{SM}}$ are given[12] by 19.5 pb for gg-fusion, 1.56 pb for VBF, 0.697 pb for $W \to Wh_{SM}$, and 0.394 pb for $Z \to Zh_{SM}$, for 125 GeV h_{SM}.

where → represents the large tanβ limit. Here A_t, A_b are the soft SUSY breaking trilinear parameters, $X_t = A_t - \mu\cot\beta$, and h_t, h_b are Yukawa coupling parameters. X_t also appears in the off-diagonal element of the scalar top mass-squared matrix. The $-\epsilon$ is the value of $-(m_A^2 + M_Z^2)s_\beta c_\beta$ for a large tanβ. The flavor-tuned (FT) model, where $\alpha \approx 0$, requires a cancellation of this $-\epsilon$ term by Loop$_{12}$ leading to $\mathcal{M}^2_{H,12} = 0$.

The 125 GeV mass of h suggests the X_t value[13] $X_t \simeq -\sqrt{6}M_{\rm SUSY}$, however, taking $X_t \simeq -2M_{\rm SUSY}$ close to the maximal mixing is sufficient to reproduce the observed m_h for $M_{\rm SUSY} \geq 1$ TeV. If we take $\mu/M_{\rm SUSY} = 1$, $\mathcal{M}^2_{H,12} = 0$ is reproduced when $m_A = (3M_Z/2,\ M_Z,\ M_Z/2)$ corresponding, respectively, to tan$\beta = (45,\ 28,\ 17)$. Thus, small m_A and large tanβ values are implied. The masses of heavier Higgs H, charged Higgs $H^\pm$ are given in the tree-level as

$$M_H^2 = M_A^2 + M_Z^2 \sin^2 2\beta, \quad M_{H^+}^2 = M_A^2 + M_W^2 \tag{10}$$

The radiative corrections to these tree-level relations are expected to be small. All these Higgs bosons should have masses of $\mathcal{O}(M_Z)$. Their masses are estimated in three cases of

m_A .

$$\begin{array}{cccc} \tan\beta & m_A & m_H & m_{H^+} \\ 45 & 3M_Z/2 & 164\text{ GeV} & 159\text{ GeV} \\ 28 & M_Z & 129\text{ GeV} & 122\text{ GeV} \\ 14 & M_Z/2 & 102\text{ GeV} & 92\text{ GeV} \end{array} \tag{11}$$

where the value of $\tan\beta$ is obtained from the cancellation condition $\mathcal{M}^2_{H,12} = 0$ in Eq. (9): $X_t/M_{\rm SUSY} \simeq A_t/M_{\rm SUSY} \simeq -2$ and $\mu/M_{\rm SUSY} \simeq 1$ are used. The $m_A = 3M_Z/2$ case is already excluded by the CMS[14], but the lighter m_A solutions are still viable.

The other neutral CP-even Higgs $H = H^0_d$ couples to the $bb, \tau\tau, \mu\mu$ channels. The $\tau\tau$ channel provides a way to detect H and A [1]. The XC and R^H results from the couplings[3] :

$$r^H_{VV} = c_{\beta-\alpha},\ r^H_{tt} = r^H_{cc} = \frac{s_\alpha}{s_\beta},\ r^H_{bb} = r^H_{\tau\tau} = r^H_{\mu\mu} = \frac{c_\alpha}{c_\beta},\ r^H_{gg} = \frac{1.03\frac{s_\alpha}{s_\beta} + (-0.059 + i0.081)\frac{c_\alpha}{c_\beta}}{1.03 + (-0.059 + i0.081)},$$
$$r^H_{\gamma\gamma} = \frac{\frac{7}{4}1.19\ \cos(\beta-\alpha) - \frac{4}{9}1.03\frac{s_\alpha}{s_\beta} - \frac{1}{9}(-0.059 + i0.081)\frac{c_\alpha}{c_\beta}}{\frac{7}{4}1.19 - \frac{4}{9}1.03 - \frac{1}{9}(-0.059 + i0.081)} \tag{12}$$

where we use the same numbers of triangle functions as Eqs. (4),(5) by simply taking $m_H = m_h = 125$ GeV. Unlike h, the H decays to b-quark pairs are dominant.

The XC and R^A formulae for CP-odd neutral state A are obtained from the couplings :

$$r^A_{VV} = 0,\ r^A_{tt} = r^A_{cc} = -i\gamma_5\ t_\beta^{-1},\ r^H_{bb} = r^H_{\tau\tau} = r^H_{\mu\mu} = -i\gamma_5\ t_\beta.$$
$$r^A_{gg} = \frac{1.57 t_\beta^{-1} + (-0.066 + i0.080)t_\beta}{1.03 + (-0.059 + i0.081)},$$
$$r^A_{\gamma\gamma} = \frac{-\frac{4}{9}1.57 t_\beta^{-1} - \frac{1}{9}(-0.066 + i0.080)t_\beta}{\frac{7}{4}1.19 - \frac{4}{9}1.03 - \frac{1}{9}(-0.059 + i0.081)} \tag{13}$$

where we again simply take $m_A = m_h$ =125 GeV. The A boson is not coupled to VV.

By taking m_H and m_A commonly 125 GeV we obtain the following values of XC at $\tan\beta = 10$.

$$\begin{array}{cccccccc} & V\tau & qb & g\tau & g\gamma & gW & gZ & V\gamma \\ H & 0.02 & 0.02 & 1.6 & 0.0 & 0.0 & 0.0 & 0.0 \\ A & 0.0 & 0.0 & 1.45 & 0.0 & 0.0 & 0.0 & 0.0 \end{array} \tag{14}$$

where we ignore deferences in QCD K-factors of H and A production relative to h_{SM}. Here $g\tau = 1.6$ means that 1.6 times more events compared with h_{SM} with the same mass. The



FIG. 2. tanβ dependence of the cross-section ratios $g\tau$ of H, A to the SM Higgs. $g\tau$ for $H(A)$ is solid black(dashed blue).

tanβ dependence of $g\tau = gb = g\mu$ is shown in Fig. 2. When the $H = H_d^0$ or A are detected in this channel, tanβ will be determined at the same time.

For the A-state the $\mu\mu$ channel is also promising. Its collider signature is investigated in refs.[15, 16].

The decay Branching fractions(%) of H, A are given by

$$\begin{array}{ccccccc}
\alpha = 0 & bb & \tau\tau & gg & \gamma\gamma & WW^* & ZZ^* \\
H & 91 & 9 & 0.13 & 0.001 & 0.049 & 0.006 \\
A & 91 & 9 & 0.11 & 0.00 & 0 & 0
\end{array}
\qquad
\begin{array}{ccccccc}
\alpha = 0.06 & bb & \tau\tau & gg & \gamma\gamma & WW^* & ZZ^* \\
H & 91 & 9 & 0.12 & 0.019 & 0.008 & 0.001 \\
A & 91 & 9 & 0.12 & 0.000 & 0 & 0
\end{array} \tag{15}$$

They dominantly decay to $b\bar{b}$. The H^+ decays[17] dominantly to $\tau^+\nu_\tau$ and $c\bar{s}$.

The light H^+ can be probed in top decay $t \to H^+b$. The tanβ dependence of the $BF(t \to H^+b)$ is shown in Fig. 3.

The experimental upper limit[1] is $BF(t \to H^+b) < 0.05(0.01)$ for $m_{H^+} = 90(130)$ GeV, which allows 5 <tan$\beta < 11$ ($3 \leq$ tan$\beta \leq 21$) for $m_{H^+} = 125(90)$ GeV case.

The light H^+ gives a large loop-contribution to $b \to s\gamma$, but this can be cancelled by the chargino loop. A recent study[19] of natural SUSY which includes the light m_{H^+} is consistent with the the $b \to s\gamma$ experimental branching fraction.

Concluding remarks



FIG. 3. tanβ dependence of the branching fraction $BF(t \to H^+ b)$. $m_{H^+} = 125$ GeV is taken. We use $\Gamma^t_{\rm tot} = 1.3$ GeV[18].

The observed enhancement of the diphoton cross section relative to the SM Higgs prediction can be explained in the flavor-tuned model with $h = H_u^0$. The maximum enhancement reaches 3[8] in the case of no mixing $\alpha = 0$ in the (H_u^0, H_d^0)-basis. This diphoton enhancement is due to reduction of the $b\bar{b}$ decay width compared to h_{SM}. There is a corresponding reduction of the h to $\tau\tau$ signal.

The vanishing of the neutral Higgs mixing angle α requires that the masses M_H, M_A and M_H^+ are the electroweak scale. Their production and decays are unambiguously predicted by the FT model and can be probed by the LHC experiments. In reality, the value of α should not be precisely zero, since its smallness depends on a cancellation of tree-level and radiative corrections in the 12 element of the neutral Higgs mass matrix. Thus, the Higgs branching fractions could be shifted somewhat with non-zero but small α, as shown in Fig. 1 for the 125 GeV h. A non-zero alpha will also allow some latitude in the predicted H, A, H^+ mass spectrum from the idealized $\alpha = 0$ result. If one of the neutral SUSY Higgs bosons has a mass similar to that of the h-boson, there may be confusion in separating the signals. The allowed "sweet-spot" for a A-boson of mass $\sim m_h$ is tan$\beta = 5$-10.[14]

Acknowledgements

W.-Y. K. and M. I. thank the hospitality of National Center for Theoretical Sciences and Academia Sinica while this work was completed. This work was supported in part by the U.S. Department of Energy under grants No. DE-FG02-95ER40896 and DE-FG02-12ER41811, in part by KAKENHI(2274015, Grant-in-Aid for Young Scientists(B)).

---

[*] barger@pheno.wisc.edu

[†] mishida@wisc.edu

[‡] keung@uic.edu

[1] ATLAS Collaboration, https://twiki.cern.ch/twiki/bin/view/AtlasPublic

[2] CMS Collaboration, http://cdsweb.cern.ch/collection/CMS?ln=en

[3] V. Barger, H. E. Logan, G. Shaughnessy, arXiv:0902.0170.

[4] D. Chung et al., Phys. Rept. 407 (2005) 1-203. arXiv:hep-ph/0312378

[5] H. Baer and X. Tata, "Weak scale Supersymmetry: From Superfields to Scattering Events", Cambridge University Press 2006.

[6] V. Barger, M. Ishida, and W.-Y. Keung, arXiv:1203.3456[hep-ph], Phys. Rev. Lett. **108**, 261801 (2012).

[7] A. Denner, S. Heinemeyer, I. Puljak, D. Rebuzzi, and M. Spira, arXiv:1107.5909.

[8] J. Chang, K. Cheung, Po-Yan Tseng, and Tzu-Chiang Yuan, arXiv:1206.5853.

[9] M. Carena, S. Gori, N. R. Shah, C. E. Wagner, arXiv:1112.3336.

[10] H. E. Haber and R. Hempfling, arXiv: hep-ph/9307201.

[11] The TEVNPH Working Group for the CDF and D0 Collaboration, FERMILAB-CONF-12-065-E . arXiv: 1207.0449[hep-ex].

[12] LHC Physics Web. https://twiki.cern.ch/twiki/bin/view/LHCPhysics/CERNYellowReportPageAt8TeV

[13] V. Barger, P. Huang, M. Ishida, and W.-Y. Keung, arXiv:1206.1777[hep-ph].

[14] Partcle data group 2012, Review on Higgs boson, p.45.

[15] V. Barger and Chung Kao, hep-ph/9711328.

[16] S. Dawson, D. Dicus, and Chung Kao, hep-ph/0208063.

[17] V. Barger, R. J. N. Phillips, and D. P. Roy, hep-ph/9311372.

[18] V. M. Abazov et al, the D0 Collaboration, arXiv:0906.5326[hep-ex].

[19] V. Barger, J. L. Hewett, R. J. N. Phillips, Phys. Rev. D**41**, 3421 (1990).